
Whip-poor-will (Caprimulgus vociferus)

Species Assessment Scores*

| State rarity: | 2 |
|--------------------------|-----|
| State threats: | 3 |
| State population trend: | 4 |
| Global abundance: | 3 |
| Global distribution: | 3 |
| Global threats: | 3 |
| Global population trend: | 4 |
| Mean Risk Score: | 3.1 |
| Area of importance: | 3 |

^{*} Please see the <u>Description of Vertebrate Species</u>
<u>Summaries (Section 3.1.1)</u> for definitions of criteria and scores.



Ecological Landscape Associations

Please note that this is not a range map. Shading does not imply that the species is present throughout the Landscape, but represents the probability that the species occurs somewhere in the Landscape.

Landscape -community Combinations of Highest Ecological Priority

| Ecological Landscape | Community |
|---------------------------|-------------------------------|
| Central Sand Hills | Central sands pine-oak forest |
| Central Sand Hills | Southern dry forest |
| Central Sand Hills | Southern dry-mesic forest |
| Central Sand Plains | Central sands pine-oak forest |
| Central Sand Plains | Oak barrens |
| Central Sand Plains | Pine barrens |
| Central Sand Plains | Southern dry forest |
| Central Sand Plains | Southern dry-mesic forest |
| North Central Forest | Bedrock glade |
| Northeast Sands | Northern dry forest |
| Northeast Sands | Northern dry -mesic forest |
| Northeast Sands | Pine barrens |
| Northern Highland | Northern dry -mesic forest |
| Northwest Sands | Northern dry forest |
| Northwest Sands | Northern dry -mesic forest |
| Northwest Sands | Pine barrens |
| Southeast Glacial Plains | Oak woodland |
| Southeast Glacial Plains | Southern dry forest |
| Southeast Glacial Plains | Southern dry-mesic forest |
| Western Coulee and Ridges | Bedrock glade |
| Western Coulee and Ridges | Oak barrens |
| Western Coulee and Ridges | Oak woodland |
| Western Coulee and Ridges | Pine relict |
| Western Coulee and Ridges | Southern dry forest |
| Western Coulee and Ridges | Southern dry-mesic forest |

Threats and Issues

• Very little is known about population trend, distribution, habitat selection and limiting factors either in Wisconsin or in the tropical wintering grounds.

- As a ground nester, may be sensitive to predation from expanding meso-predator complexes in many areas of Wisconsin.
- Tropical deforestation may be impacting populations.
- Dependent on large moths for food. Some have speculated that agricultural chemicals may be limiting moth populations.

Priority Conservation Actions

- Demographic research to identify potential limiting factors.
- Long-term population trend survey in the Upper Midwest.
- Conserve large blocks of oak woodlands and dry forest.